

MID-TERM EXAMINATION (2017-18) SUBJECT: CHEMISTRY

CLASS:

TIME - 3 HOURS

Lovish MAXIMUM MARKS - 60

Important instructions:

- 1. All guestions are compulsory.
- 2. Q. No. 1 to 8 are very short answer type questions and carry 1 mark each.
 - 3. Q. No. 9 to 14 are short answer type questions carrying 2 marks each.
 - 4. Q. No. 15 to 24 are short answer type questions carrying 3 marks each.
 - 5. Q. No. 25 and 26 are long answer type questions carrying 5 marks each.
 - What is the difference between 5.0 g and 5.00 g? Which is more precise?
 - What is the actual value of charge and mass of an electron?
- Which of the following pairs of elements would have a more electron gain enthalpy: Will a the electronic configuration and the number of u D To T (d) N To O (s)
- 4. How many atoms of fluorine are present in 1.9 x 10⁻⁶ g of fluorine? (atomic mass of fluorine = 19 u)
 - 5. Two flasks of equal volumes contain N₂ and O₂ gases at same temperature and pressure. Which will have greater number of molecules? Justify
 - What do you mean by isolated system? Give an example.
 - If 6.3 g of NaHCO₃ are added to 15.0 g of CH₃COOH, the residue is found to weigh 18.0 g. What is the mass of CO2 released in the reaction?
 - 8. Noble gases form neither anions nor cations. Why?
 - 9. Define: (i) Bond angle and (ii) Bond length
 - Explain by giving reasons: (i) Beryllium has higher ionisation enthalpies than boron. (ii) Oxygen has lower ionisation enthalpies than nitrogen and Fluorine
 - Define isothermal and adiabatic processes.
 - 12. Name one compound each involving sp and sp² hybridisation. Draw the structure of each molecule.

- 13. What is the wavelength of the light emitted when the electron in the hydrogen atom undergoes transition from an energy level with n=4 to energy level n=2?
- 14. Calculate the internal energy change in the following case: When a system absorbs 15 kJ of heat and does 5 kJ of work.
- 15. A proton is accelerated to $1/10^{th}$ of the velocity of light. If its velocity can be measured with a precision of 0.5%, what must be its uncertainty in position? (mass of proton = 1.66×10^{-27} kg.)
- 16. What is the concentration of sucrose (C₁₂H₂₂O₁₁) in moles/L if 40 g of sugar are dissolved in enough water to make a final volume up to 2 L?

have same Quantum no.

Mid-term/XI/ Chemistry/25.09.2017 1 | Page

- 17. A flask was heated from 27°C to 227°C at constant pressure. Calculate the volume of the flask if 0.1 dm³ of air measured at 227°C was expelled from the flask

 18. State Boyle's law and show how it can be explained by kinetic theory of gases? How can the law be verified graphically?

 OR

 (a) Define Charle's law.
 - (b) A certain amount of gas occupies a volume of 400 ml at 17°C. To what temperature should it be heated so that the volume is reduced to half.
 - 19. Write the electronic configuration of the following: (i) O2- (ii) Ca (iii) Cr
 - 20. An Oxide of nitrogen has the percentage composition N= 25.94 and O = 74.06. Calculate the empirical formula of the compound.
 - 21. The kinetic energy of an electron is 3.0 x 10⁻²⁵ J. Calculate its wavelength.
 - 22. (a) Define Pauli's exclusion principle. It is a private of the private of the
 - (b) Write the electronic configuration and the number of unpaired electrons in Fe²⁺ ion
 - Write three important postulates of kinetic theory of gases. What are the two postulates of theory are defective .Explain.

OR

What are the different types of molecular speed? Explain them by giving mathematical

- 24. Define ionisation enthalpy. List the various factors on which the ionisation enthalpy depends.
- 25. (a) With the help of molecular orbital theory explain, why hydrogen forms di atomic molecule while helium remains mono atomic.
 - (b) On the basis of hybridisation discuss and draw the shape of the following molecules: (i) H₂O (ii) C₂H₂
- 26. (i) What is the relationship between wavelength and momentum of a particle?
 - (ii) Which element has only one electron in the d orbital?
 - (iii) The work function of Cs atom is 1.9 eV. Calculate (a) threshold wavelength and (b) the threshold frequency of the radiation. If the caesium element is irradiated with a radiation of wavelength 500 nm. Calculate the kinetic energy and the velocity of the ejected photo electrons.

- (a) Which of the following represent ground state configurations and which are excited state configurations: (i) 1s²2s²2p⁴ (ii) 1s²2s²2p⁶3s¹3p¹ (iii) 1s²2s²2p⁶3s²3p⁴
- (b) How many unpaired electrons are present in Mn²⁺
- Find the total number of electrons in a molecule of phosphoric acid H_3PO_4 (atomic numbers of H = 1, P = 15, O = 96%)
- (d) How many protons are present in 5.6 L of oxygen at NTP, using O-16 isotope only

- (a) Which of the following represent ground state configurations and which are excited state configurations: (i) 1s²2s²2p⁴ (ii) 1s²2s²2p⁶3s¹3p¹ (iii) 1s²2s²2p⁶3s²3p⁴
- (b) How many unpaired electrons are present in Mn²⁺
- Find the total number of electrons in a molecule of phosphoric acid H_3PO_4 (atomic numbers of H = 1, P = 15, O = \P
- (d) How many protons are present in 5.6 L of oxygen at NTP, using O-16 isotope only

Mid-term/XI/ Chemistry/25.09.2017 2 | Page