

NCERT Solutions for Class 12 Chemistry Part 1 Chapter 7

The p-Block Elements Class 12

Chapter 7 The p-Block Elements Exercise Solutions

In text : Solutions of Questions on Page Number : 169

Q1 :

Why are pentahalides more covalent than trihalides?

Answer :

In pentahalides, the oxidation state is +5 and in trihalides, the oxidation state is +3. Since the metal ion with a high charge has more polarizing power, pentahalides are more covalent than trihalides.

Q2 :

Why is BiH₃ the strongest reducing agent amongst all the hydrides of Group 15 elements?

Answer :

As we move down a group, the atomic size increases and the stability of the hydrides of group 15 elements decreases. Since the stability of hydrides decreases on moving from NH₃ to BiH₃, the reducing character of the hydrides increases on moving from NH₃ to BiH₃.

Q3 :

Why is N₂ less reactive at room temperature?

Answer :

The two N atoms in N₂ are bonded to each other by very strong triple covalent bonds. The bond dissociation energy of this bond is very high. As a result, N₂ is less reactive at room temperature.


Q4 :

Mention the conditions required to maximise the yield of ammonia.

Answer :

Ammonia is prepared using the Haber's process. The yield of ammonia can be maximized under the following conditions:

(i) High pressure ($\propto \frac{1}{200}$ atm)




Q8 :

What happens when white phosphorus is heated with concentrated NaOH solution in an inert atmosphere of CO_2 ?

Answer :

White phosphorous dissolves in boiling NaOH solution (in a CO_2 atmosphere) to give phosphine, PH_3 .




Q9 :

What happens when PCl_5 is heated?

Answer :


All the bonds that are present in PCl_5 are not similar. It has three equatorial and two axial bonds. The equatorial bonds are stronger than the axial ones. Therefore, when PCl_5 is heated strongly, it decomposes to form PCl_3 .



Q10 :

Write a balanced equation for the hydrolytic reaction of PCl_5 in heavy water.

Answer :




Q11 :

What is the basicity of H_3PO_4 ?

Answer :

H_3PO_4




Since there are three OH groups present in H_3PO_4 , its basicity is three i.e., it is a tribasic acid.

Q12 :

What happens when H_3PO_3 is heated?

Answer :

H_3PO_3 , on heating, undergoes disproportionation reaction to form PH_3 and H_3PO_4 . The oxidation numbers of P in H_3PO_3 , PH_3 , and H_3PO_4 are +3, -3, and +5 respectively. As the oxidation number of the same element is decreasing and increasing during a particular reaction, the reaction is a disproportionation reaction.



Q13 :

List the important sources of sulphur.

Answer :

Sulphur mainly exists in combined form in the earth's crust primarily as sulphates [gypsum ($\text{CaSO}_4 \cdot 2\text{H}_2\text{O}$), Epsom salt ($\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$), baryte (BaSO_4)] and sulphides [(galena (PbS), zinc blends (ZnS), copper pyrites (CuFeS_2)].


Q14 :

Write the order of thermal stability of the hydrides of Group 16 elements.

Answer :

The thermal stability of hydrides decreases on moving down the group. This is due to a decrease in the bond dissociation enthalpy (H-E) of hydrides on moving down the group.

Therefore,



Q15 :

Why is H₂O a liquid and H₂S a gas?

Answer :

H₂O has oxygen as the central atom. Oxygen has smaller size and higher electronegativity as compared to sulphur. Therefore, there is extensive hydrogen bonding in H₂O, which is absent in H₂S. Molecules of H₂S are held together only by weak van der Waal's forces of attraction.

Hence, H₂O exists as a liquid while H₂S as a gas.

Q16 :

Which of the following does not react with oxygen directly?


Zn, Ti, Pt, Fe

Answer :



Pt is a noble metal and does not react very easily. All other elements, Zn, Ti, Fe, are quite reactive. Hence, oxygen does not react with platinum (Pt) directly.

Q17 :

Complete the following reactions:



Answer :




Q18 :

Why does O₃ act as a powerful oxidising agent?

Answer :

Ozone is not a very stable compound under normal conditions and decomposes readily on heating to give a molecule of oxygen and nascent oxygen. Nascent oxygen, being a free radical, is very reactive.




Therefore, ozone acts as a powerful oxidising agent.

Q19 :

How is O₃ estimated quantitatively?

Answer :

Quantitatively, ozone can be estimated with the help of potassium iodide. When ozone is made to react with potassium iodide solution buffered with a borate buffer (pH 9.2), iodine is liberated. This liberated iodine can be titrated against a standard solution of sodium thiosulphate using starch as an indicator. The reactions involved in the process are given below.



Q20 :

What happens when sulphur dioxide is passed through an aqueous solution of Fe(III) salt?

Answer :

SO₂ acts as a reducing agent when passed through an aqueous solution containing Fe(III) salt. It reduces Fe(III) to Fe(II) i.e., ferric ions to ferrous ions.




Q21 :

Comment on the nature of two S-O bonds formed in SO₂ molecule. Are the two S-O bonds in this molecule equal?

Answer :

The electronic configuration of S is 1s² 2s² 2p⁶ 3s² 3p⁴.

During the formation of SO₂, one electron from 3p orbital goes to the 3d orbital and S undergoes sp² hybridization. Two of these orbitals form sigma bonds with two oxygen atoms and the third contains a lone pair. p-orbital and d-orbital contain an unpaired electron each. One of these electrons forms p-d bond with one oxygen atom and the other forms p-d bond with the other oxygen. This is the reason SO₂ has a bent structure. Also, it is a resonance hybrid of structures I and II.



Both S-O bonds are equal in length (143 pm) and have a multiple bond character.

Q22 :

How is the presence of SO₂ detected?

Answer :

SO₂ is a colourless and pungent smelling gas.

It can be detected with the help of potassium permanganate solution. When SO₂ is passed through an acidified potassium permanganate solution, it decolorizes the solution as it reduces MnO₄⁻ ions to Mn²⁺ ions.



Q23 :

Mention three areas in which H₂SO₄ plays an important role.

Answer :

Sulphuric acid is an important industrial chemical and is used for a lot of purposes. Some important uses of sulphuric acid are given below.

(i) It is used in fertilizer industry. It is used to make various fertilizers such as ammonium sulphate and calcium super phosphate.

(ii) It is used in the manufacture of pigments, paints, and detergents.

(iii) It is used in the manufacture of storage batteries.

Q24 :

Write the conditions to maximize the yield of H_2SO_4 by Contact process.

Answer :

Manufacture of sulphuric acid by Contact process involves three steps.

1. Burning of ores to form SO_2
2. Conversion of SO_2 to SO_3 by the reaction of the former with O_2

(V_2O_5 is used in this process as a catalyst.)


3. Absorption of SO_3 in H_2SO_4 to give oleum ($H_2S_2O_7$)

The key step in this process is the second step. In this step, two moles of gaseous reactants combine to give one mole of gaseous product. Also, this reaction is exothermic. Thus, in accordance with Le Chatelier's principle, to obtain the maximum amount of SO_3 gas, temperature should be low and pressure should be high.

Q25 :

Why is $K_{a_2} \ll K_{a_1}$ for H_2SO_4 in water?

Answer :



It can be noticed that $K_{a_1} \gg K_{a_2}$

This is because a neutral H_2SO_4 has a much higher tendency to lose a proton than the negatively charged HSO_4^- . Thus, the former is a much stronger acid than the latter.

Q26 :

Considering the parameters such as bond dissociation enthalpy, electron gain enthalpy and hydration enthalpy, compare the oxidising power of F_2 and Cl_2 .

Answer :

Fluorine is a much stronger oxidizing agent than chlorine. The oxidizing power depends on three factors.

1. Bond dissociation energy
2. Electron gain enthalpy
3. Hydration enthalpy

The electron gain enthalpy of chlorine is more negative than that of fluorine. However, the bond dissociation energy of fluorine is much lesser than that of chlorine. Also, because of its small size, the hydration energy of fluorine is much higher than that of chlorine. Therefore, the latter two factors more than compensate for the less negative electron gain enthalpy of fluorine. Thus, fluorine is a much stronger oxidizing agent than chlorine.

Q27 :

Give two examples to show the anomalous behaviour of fluorine.

Answer :

Anomalous behaviour of fluorine

- (i) It forms only one oxoacid as compared to other halogens that form a number of oxoacids.
- (ii) Ionisation enthalpy, electronegativity, and electrode potential of fluorine are much higher than expected.

Q28 :

Sea is the greatest source of some halogens. Comment.

Answer :

Sea water contains chlorides, bromides, and iodides of Na, K, Mg, and Ca. However, it primarily contains NaCl. The deposits of dried up sea beds contain sodium chloride and carnallite, $KCl \cdot MgCl_2 \cdot 6H_2O$. Marine life also contains iodine in their systems. For example, sea weeds contain upto 0.5% iodine as sodium iodide. Thus, sea is the greatest source of halogens.

Q29 :

Give the reason for bleaching action of Cl_2 .

Answer :

When chlorine reacts with water, it produces nascent oxygen. This nascent oxygen then combines with the coloured substances present in the organic matter to oxidize them into colourless substances.



Coloured substances + [O] → Oxidized colourless substance

Q30 :

Name two poisonous gases which can be prepared from chlorine gas.

Answer :

Two poisonous gases that can be prepared from chlorine gas are

(i) Phosgene (COCl_2)

(ii) Mustard gas ($\text{ClCH}_2\text{CH}_2\text{SCH}_2\text{CH}_2\text{Cl}$)

Q31 :

Why is ICl more reactive than I_2 ?

Answer :

ICl is more reactive than I_2 because I-Cl bond in ICl is weaker than I-I bond in I_2 .

Q32 :

Why is helium used in diving apparatus?

Answer :


Air contains a large amount of nitrogen and the solubility of gases in liquids increases with increase in pressure. When sea divers dive deep into the sea, large amount of nitrogen dissolves in their blood. When they come back to the surface, solubility of nitrogen decreases and it separates from the blood and forms small air bubbles. This leads to a dangerous medical condition called bends. Therefore, air in oxygen cylinders used for diving is diluted with helium gas. This is done as He is sparingly less soluble in blood.

Q33 :

Balance the following equation: $\text{XeF}_6 + \text{H}_2\text{O} \rightarrow \text{XeO}_2\text{F}_2 + \text{HF}$

Answer :

Balanced equation



Q34 :

Why has it been difficult to study the chemistry of radon?

Answer :

It is difficult to study the chemistry of radon because it is a radioactive substance having a half-life of only 3.82 days. Also, compounds of radon such as RnF_2 have not been isolated. They have only been identified.

<< Previous Chapter 6 : General Principles and Processes of Isolation of Elements Next Chapter 8 : The d-and f-Block Elements



Exercise : Solutions of Questions on Page Number : 207

Q1 :

Discuss the general characteristics of Group 15 elements with reference to their electronic configuration, oxidation state, atomic size, ionisation enthalpy and electronegativity.

Answer :

General trends in group 15 elements

(i) Electronic configuration: All the elements in group 15 have 5 valence electrons. Their general electronic configuration is $ns^2 np^3$.

(ii) Oxidation states: All these elements have 5 valence electrons and require three more electrons to complete their octets. However, gaining electrons is very difficult as the nucleus will have to attract three more electrons. This can take place only with nitrogen as it is the smallest in size and the distance between the nucleus and the valence shell is relatively small. The remaining elements of this group show a formal oxidation state of -3 in their covalent compounds. In addition to the -3 state, N and P also show -1 and -2 oxidation states.

All the elements present in this group show +3 and +5 oxidation states. However, the stability of +5 oxidation state decreases down a group, whereas the stability of +3 oxidation state increases. This happens because of the inert pair effect.

(iii) Ionization energy and electronegativity

First ionization decreases on moving down a group. This is because of increasing atomic sizes. As we move down a group, electronegativity decreases, owing to an increase in size.

(iv) Atomic size: On moving down a group, the atomic size increases. This increase in the atomic size is attributed to an increase in the number of shells.

Q2 :

Why does the reactivity of nitrogen differ from phosphorus?

Answer :

Nitrogen is chemically less reactive. This is because of the high stability of its molecule, N_2 . In N_2 , the two nitrogen atoms form a triple bond. This triple bond has very high bond strength, which is very difficult to break. It is because of nitrogen's small size that it is able to form $p\text{-}p$ bonds with itself. This property is not exhibited by atoms such as phosphorus. Thus, phosphorus is more reactive than nitrogen.

Q3 :

Discuss the trends in chemical reactivity of group 15 elements.

Answer :

General trends in chemical properties of group - 15

(i) Reactivity towards hydrogen:

The elements of group 15 react with hydrogen to form hydrides of type EH_3 , where $E = N, P, As, Sb, \text{ or } Bi$. The stability of hydrides decreases on moving down from NH_3 to BiH_3 .

(ii) Reactivity towards oxygen:

The elements of group 15 form two types of oxides: E_2O_3 and E_2O_5 , where $E = N, P, As, Sb, \text{ or } Bi$. The oxide with the element in the higher oxidation state is more acidic than the other. However, the acidic character decreases on moving down a group.

(iii) Reactivity towards halogens:

The group 15 elements react with halogens to form two series of salts: EX_3 and EX_5 . However, nitrogen does not form NX_5 as it lacks the d -orbital. All trihalides (except NX_3) are stable.

(iv) Reactivity towards metals:

The group 15 elements react with metals to form binary compounds in which metals exhibit -3 oxidation states.

Q4 :

Why does NH_3 form hydrogen bond but PH_3 does not?

Answer :

Nitrogen is highly electronegative as compared to phosphorus. This causes a greater attraction of electrons towards nitrogen in NH_3 than towards phosphorus in PH_3 . Hence, the extent of hydrogen bonding in PH_3 is very less as compared to NH_3 .

Q5 :

How is nitrogen prepared in the laboratory? Write the chemical equations of the reactions involved.

Answer :

An aqueous solution of ammonium chloride is treated with sodium nitrite.



NO and HNO₃ are produced in small amounts. These are impurities that can be removed on passing nitrogen gas through aqueous sulphuric acid, containing potassium dichromate.

Q6 :

How is ammonia manufactured industrially?

Answer :

Ammonia is prepared on a large-scale by the Haber's process.




Q7 :

Illustrate how copper metal can give different products on reaction with HNO₃.

Answer :

Concentrated nitric acid is a strong oxidizing agent. It is used for oxidizing most metals. The products of oxidation depend on the concentration of the acid, temperature, and also on the material undergoing oxidation.




Q8 :


Give the resonating structures of NO₂ and N₂O₅.

Answer :

(1)



(2)



Q9 :

The HNH angle value is higher than HPH, HAsH and HSbH angles. Why? [Hint: Can be explained on the basis of sp^3 hybridisation in NH_3 and only $s-p$ bonding between hydrogen and other elements of the group].

Answer :

Hydride NH_3 , PH_3 , AsH_3 , SbH_3

H-M-H angle 107° , 92° , 91° , 90°

The above trend in the H-M-H bond angle can be explained on the basis of the electronegativity of the central atom. Since nitrogen is highly electronegative, there is high electron density around nitrogen. This causes greater repulsion between the electron pairs around nitrogen, resulting in maximum bond angle. We know that electronegativity decreases on moving down a group. Consequently, the repulsive interactions between the electron pairs decrease, thereby decreasing the H-M-H bond angle.

Q10 :

Why does $R_3P=O$ exist but $R_3N=O$ does not (R = alkyl group)?

Answer :

N (unlike P) lacks the d -orbital. This restricts nitrogen to expand its coordination number beyond four. Hence, $R_3N=O$ does not exist.

Q11 :

Explain why NH_3 is basic while BiH_3 is only feebly basic.

Answer :

NH_3 is distinctly basic while BiH_3 is feebly basic.

Nitrogen has a small size due to which the lone pair of electrons is concentrated in a small region. This means that the charge density per unit volume is high. On moving down a group, the size of the central atom increases and the charge gets distributed over a large area decreasing the electron density. Hence, the electron donating capacity of group 15 element hydrides decreases on moving down the group.

Q12 :

Nitrogen exists as diatomic molecule and phosphorus as P₄. Why?



Answer :

Nitrogen owing to its small size has a tendency to form multiple bonds with itself. Nitrogen thus forms a very stable diatomic molecule, N₂. On moving down a group, the tendency to form multiple bonds decreases (because of the large size of heavier elements). Therefore, phosphorus (like other heavier metals) exists in the P₄ state.

Q13 :

Write main differences between the properties of white phosphorus and red phosphorus.

Answer :

White phosphorus	Red Phosphorus
It is a soft and waxy solid. It possesses a garlic smell.	It is a hard and crystalline solid, without any smell.
It is poisonous.	It is non-poisonous.
It is insoluble in water but soluble in carbon disulphide.	It is insoluble in both water and carbon disulphide.
It undergoes spontaneous combustion in air.	It is relatively less reactive.
In both solid and vapour states, it exists as a P ₄ molecule. 	It exists as a chain of tetrahedral P ₄ units. 

Q14 :

Why does nitrogen show catenation properties less than phosphorus?

Answer :


Catenation is much more common in phosphorous compounds than in nitrogen compounds. This is because of the relative weakness of the N-N single bond as compared to the P-P single bond. Since nitrogen atom is smaller, there is greater repulsion of electron density of two nitrogen atoms, thereby weakening the N-N single bond.

Q15 :

Give the disproportionation reaction of H_3PO_3 .

Answer :

On heating, orthophosphorus acid (H_3PO_3) disproportionates to give orthophosphoric acid (H_3PO_4) and phosphine (PH_3). The oxidation states of P in various species involved in the reaction are mentioned below.



Q16 :

Can PCl_5 act as an oxidising as well as a reducing agent? Justify.

Answer :

PCl_5 can only act as an oxidizing agent. The highest oxidation state that P can show is +5. In PCl_5 , phosphorus is in its highest oxidation state (+5). However, it can decrease its oxidation state and act as a reducing agent.

Q17 :

Justify the placement of O, S, Se, Te and Po in the same group of the periodic table in terms of electronic configuration, oxidation state and hydride formation.

Answer :

The elements of group 16 are collectively called chalcogens.

(i) Elements of group 16 have six valence electrons each. The general electronic configuration of these elements is ns^2np^4 , where n varies from 2 to 6.

(ii) **Oxidation state:**

As these elements have six valence electrons ($ns^2 np^4$), they should display an oxidation state of -2. However, only oxygen predominantly shows the oxidation state of -2 owing to its high electronegativity. It also exhibits the oxidation state of -1 (H_2O_2), zero (O_2), and +2 (OF_2). However, the stability of the -2 oxidation state decreases on moving down a group due to a decrease in the electronegativity of the elements. The heavier elements of the group show an oxidation state of +2, +4, and +6 due to the availability of d -orbitals.

(iii) Formation of hydrides:

These elements form hydrides of formula H_2E , where $E = O, S, Se, Te, PO$. Oxygen and sulphur also form hydrides of type H_2E_2 . These hydrides are quite volatile in nature.

Q18 :

Why is dioxygen a gas but sulphur a solid?

Answer :

Oxygen is smaller in size as compared to sulphur. Due to its smaller size, it can effectively form $p\text{-}p$ bonds and form $O_2(O=O)$ molecule. Also, the intermolecular forces in oxygen are weak van der Waals, which cause it to exist as gas. On the other hand, sulphur does not form M_2 molecule but exists as a puckered structure held together by strong covalent bonds. Hence, it is a solid.

Q19 :

Knowing the electron gain enthalpy values for $O \rightarrow O^-$ and $O \rightarrow O^{2-}$ as -141

and 702 kJ mol^{-1} respectively, how can you account for the formation of a

large number of oxides having O^{2-} species and not O^- ?

(Hint: Consider lattice energy factor in the formation of compounds).

Answer :

Stability of an ionic compound depends on its lattice energy. More the lattice energy of a compound, more stable it will be.

Lattice energy is directly proportional to the charge carried by an ion. When a metal combines with oxygen, the lattice energy of the oxide involving O^{2-} ion is much more than the oxide involving O^- ion. Hence, the oxide having O^{2-} ions are more stable than oxides having O^- . Hence, we can say that formation of O^{2-} is energetically more favourable than formation of O^- .

Q20 :

Which aerosols deplete ozone?

Answer :

Freons or chlorofluorocarbons (CFCs) are aerosols that accelerate the depletion of ozone. In the presence of ultraviolet radiations, molecules of CFCs break down to form chlorine-free radicals that combine with ozone to form oxygen.

Q21 :

Describe the manufacture of H₂SO₄ by contact process?

Answer :


Sulphuric acid is manufactured by the contact process. It involves the following steps:

Step (i):

Sulphur or sulphide ores are burnt in air to form SO₂.


Step (ii):

By a reaction with oxygen, SO₂ is converted into SO₃ in the presence of V₂O₅ as a catalyst.



Step (iii):

SO₃ produced is absorbed on H₂SO₄ to give H₂S₂O₇ (oleum).



This oleum is then diluted to obtain H₂SO₄ of the desired concentration.

In practice, the plant is operated at 2 bar (pressure) and 720 K (temperature). The sulphuric acid thus obtained is 96-98% pure.

Q22 :

How is SO₂ an air pollutant?

Answer :

Sulphur dioxide causes harm to the environment in many ways:

1. It combines with water vapour present in the atmosphere to form sulphuric acid. This causes acid rain. Acid rain damages soil, plants, and buildings, especially those made of marble.
2. Even in very low concentrations, SO₂ causes irritation in the respiratory tract. It causes throat and eye irritation and can also affect the larynx to cause breathlessness.
3. It is extremely harmful to plants. Plants exposed to sulphur dioxide for a long time lose colour from their leaves. This condition is known as chlorosis. This happens because the formation of chlorophyll is affected by the presence of sulphur dioxide.

Q23 :

Why are halogens strong oxidising agents?

Answer :

The general electronic configuration of halogens is np^5 , where $n = 2-6$. Thus, halogens need only one more electron to complete their octet and to attain the stable noble gas configuration. Also, halogens are highly electronegative with low dissociation energies and high negative electron gain enthalpies. Therefore, they have a high tendency to gain an electron. Hence, they act as strong oxidizing agents.

Q24 :

Explain why fluorine forms only one oxoacid, HOF.

Answer :

Fluorine forms only one oxoacid i.e., HOF because of its high electronegativity and small size.

Q25 :

Explain why inspite of nearly the same electronegativity, oxygen forms hydrogen bonding while chlorine does not.

Answer :

Both chlorine and oxygen have almost the same electronegativity values, but chlorine rarely forms hydrogen bonding. This is because in comparison to chlorine, oxygen has a smaller size and as a result, a higher electron density per unit volume.

Q26 :

Write two uses of ClO_2 .

Answer :

Uses of ClO_2 :

- (i) It is used for purifying water.
- (ii) It is used as a bleaching agent.

Q27 :

Why are halogens coloured?

Answer :

Almost all halogens are coloured. This is because halogens absorb radiations in the visible region. This results in the excitation of valence electrons to a higher energy region. Since the amount of energy required for excitation differs for each halogen, each halogen displays a different colour.

Q28 :

Write the reactions of F₂ and Cl₂ with water.

Answer :




Q29 :

How can you prepare Cl₂ from HCl and HCl from Cl₂? Write reactions only.

Answer :

(i) Cl₂ can be prepared from HCl by Deacon's process.



(ii) HCl can be prepared from Cl₂ on treating it with water.



Q30 :

What inspired N. Bartlett for carrying out reaction between Xe and PtF₆?

Answer :

Neil Bartlett initially carried out a reaction between oxygen and PtF₆. This resulted in the formation of a red compound, $\text{O}_2^+[\text{PtF}_6]^-$.

Later, he realized that the first ionization energy of oxygen (1175 kJ/mol) and Xe (1170 kJ/mol) is almost the same. Thus, he tried to prepare a compound with Xe and PtF₆. He was successful and a red-coloured

compound, $\text{Xe}^+[\text{PtF}_6]^-$ was formed.

Q31 :

What are the oxidation states of phosphorus in the following:

(i) H_3PO_3 (ii) PCl_3 (iii) Ca_3P_2

(iv) Na_3PO_4 (v) POF_3 ?

Answer :

Let the oxidation state of p be x .

(i) H_3PO_3

$$3 + x + 3(-2) = 0$$

$$3 + x - 6 = 0$$

$$x - 3 = 0$$

$$x = +3$$

(ii) PCl_3

$$x + 3(-1) = 0$$

$$x - 3 = 0$$

$$x = +3$$

(iii) Ca_3P_2

$$3(+2) + 2(x) = 0$$

$$6 + 2x = 0$$

$$2x = -6$$

$$x = -3$$

(iv) Na_3PO_4

$$3(+1) + x + 4(-2) = 0$$

$$3 + x - 8 = 0$$

$$x - 5 = 0$$

$$x = +5$$

(v) POF_3

$$x + (-2) + 3(-1) = 0$$

$$x - 5 = 0$$

$$x = +5$$


Q32 :

Write balanced equations for the following:

(i) NaCl is heated with sulphuric acid in the presence of MnO_2 .

(ii) Chlorine gas is passed into a solution of NaI in water.

Answer :




Q33 :

How are xenon fluorides XeF_2 , XeF_4 and XeF_6 obtained?

Answer :

XeF_2 , XeF_4 and XeF_6 are obtained by a direct reaction between Xe and F_2 . The condition under which the reaction is carried out determines the product.



Q34 :

With what neutral molecule is ClO^- isoelectronic? Is that molecule a Lewis base?

Answer :

ClO^- is isoelectronic to ClF . Also, both species contain 26 electrons in all as shown.

Total electrons $\text{ClO}^- = 17 + 8 + 1 = 26$

In $\text{ClF} = 17 + 9 = 26$


ClF acts like a Lewis base as it accepts electrons from F to form ClF_3 .

Q35 :

How are XeO_3 and XeOF_4 prepared?

Answer :

(i) XeO_3 can be prepared in two ways as shown.



(ii) XeOF_4 can be prepared using XeF_6 .



Q36 :

Arrange the following in the order of property indicated for each set:

(i) F_2 , Cl_2 , Br_2 , I_2 - increasing bond dissociation enthalpy.

(ii) HF , HCl , HBr , HI - increasing acid strength.

(iii) NH_3 , PH_3 , AsH_3 , SbH_3 , BiH_3 - increasing base strength.

Answer :

(i) Bond dissociation energy usually decreases on moving down a group as the atomic size increases. However, the bond dissociation energy of F_2 is lower than that of Cl_2 and Br_2 . This is due to the small atomic size of fluorine. Thus, the increasing order for bond dissociation energy among halogens is as follows:



(ii) $\text{HF} < \text{HCl} < \text{HBr} < \text{HI}$

The bond dissociation energy of H-X molecules where $\text{X} = \text{F}, \text{Cl}, \text{Br}, \text{I}$, decreases with an increase in the atomic size. Since H-I bond is the weakest, HI is the strongest acid.

(iii) $\text{BiH}_3 < \text{SbH}_3 < \text{AsH}_3 < \text{PH}_3 < \text{NH}_3$

On moving from nitrogen to bismuth, the size of the atom increases while the electron density on the atom decreases. Thus, the basic strength decreases.

Q37 :

Which one of the following does not exist?

(i) XeOF_4 (ii) NeF_2




(iii) XeF_2 (iv) XeF_6

Answer :

NeF_2 does not exist.

Q38 :


Give the formula and describe the structure of a noble gas species which is isostructural with:



Answer :


(i)

XeF_4 is isoelectronic with ICl_4^- and has square planar geometry.




(ii)

XeF_2 is isoelectronic to IBr_2^- and has a linear structure.



(iii)

XeO_3 is isostructural to BrO_3^- and has a pyramidal molecular structure.



Q39 :

Why do noble gases have comparatively large atomic sizes?

Answer :

Noble gases do not form molecules. In case of noble gases, the atomic radii corresponds to van der Waal's radii. On the other hand, the atomic radii of other elements correspond to their covalent radii. By definition, van der Waal's radii are larger than covalent radii. It is for this reason that noble gases are very large in size as compared to other atoms belonging to the same period.

Q40 :

List the uses of Neon and argon gases.

Answer :

Uses of neon gas:

- (i) It is mixed with helium to protect electrical equipments from high voltage.
- (ii) It is filled in discharge tubes with characteristic colours.
- (iii) It is used in beacon lights.

Uses of Argon gas:

- (i) Argon along with nitrogen is used in gas-filled electric lamps. This is because Ar is more inert than N.
- (ii) It is usually used to provide an inert temperature in a high metallurgical process.
- (iii) It is also used in laboratories to handle air-sensitive substances.